

A-36

2-Magnon Peak in $\text{La}_{2-x}\text{Sr}_x\text{CuO}_4$ Observed with RIXS at the Cu K-edge

D. S. Ellis¹, Jungho Kim¹, J. P. Hill², S. Wakimoto³, R. J. Birgeneau⁴, Y. Shvyd'ko⁵, D. Casa⁵, T. Gog⁵, K. Ishii⁶, K. Ikeuchi⁶, A. Paramakanti¹, and Young-June Kim¹

¹Department of Physics, University of Toronto, Toronto, Ontario, Canada M5S 1A7

²Department of Condensed Matter Physics and Materials Science, Brookhaven National Laboratory, Upton, NY 11973

³Quantum Beam Science Directorate, Japan Atomic Energy Agency, Tokai, Ibaraki 319-1195, Japan

⁴Department of Physics, University of California, Berkeley, CA 94720-7300

⁵XOR, Advanced Photon Source, Argonne National Laboratory, Argonne, IL 60439

⁶Synchrotron Radiation Research Center, Japan Atomic Energy Agency, Hyogo 679-5148, Japan

We present a comprehensive study of the temperature and doping dependence of the 500 meV peak observed at $q=(\pi 0)$ in resonant inelastic x-ray scattering (RIXS) experiments on La_2CuO_4 . The intensity of this peak persists above the Néel temperature ($T_N=320\text{K}$), but decreases gradually with increasing temperature, reaching zero at around $T=500\text{K}$. The peak energy decreases with temperature in close quantitative accord with the behavior of the two-magnon B1g Raman peak in La_2CuO_4 and with suitable rescaling, agrees with the Raman peak shifts in $\text{EuBa}_2\text{Cu}_3\text{O}_6$ and K_2NiF_4 . The overall dispersion of this excitation in the Brillouin zone is found to be in agreement with theoretical calculations for a two-magnon excitation. Upon doping, the peak intensity decreases analogous to the Raman mode intensity and appears to track the doping dependence of the spin-correlation length. Taken together, these observations strongly suggest that the 500 meV mode is magnetic in character and is likely a two-magnon excitation.